

Cevac IBD L

COMPOSITION

CEVAC® IBD L contains the Winterfield 2512 strain of Infectious Bursal Disease virus in live, freeze dried form. The embryonated hen eggs used in the production of the vaccine are obtained from specific-pathogen-free (SPF) flocks.

INDICATIONS

For the active immunisation of healthy chickens against the disease caused by classical and very virulent strains of Infectious Bursal Disease (Gumboro Disease).

ADMINISTRATION AND DOSAGE

CEVAC® IBD L must be administered **by drinking water method**. Broilers should be vaccinated with **CEVAC® IBD L** from 10 to 18 days of age, depending on the level of maternally derived antibodies. The exact date of vaccination can be determined by checking the level of antibodies by serological methods.

STORAGE

- Store vaccine between +2°C and +8°C or 35°F and 45°F
- Protect from light

PACKAGED

- 1,000 - 2,500 and 5,000 dose vials
- 20 x 1,000 dose vials / box
- 20 x 2,500 dose vials / box
- 20 x 5,000 dose vials / box

If you wish to contact Ceva please visit their [contact page](#)

GENERAL REMINDER:

Product indications, usage instructions & withdrawal periods may vary by Country.
Always follow label instructions and consult your veterinarian or poultry health adviser.

This page contains information on veterinary pharmaceutical and biological products that are sold in several different countries and areas where they may be marketed under different trade names and pursuant to different regulatory approvals. Accordingly, ThePoultrySite and CEVA SANTE ANIMALE give no guarantee that the details presented are correct with respect to all locations.

Advertisements and/or reference to commercial products or trade names within information provided by ThePoultrySite.com does not constitute an endorsement by ThePoultrySite.com and does not imply discrimination against any other similar products. This information is aimed at veterinary surgeons, veterinary nurses, pharmacists or professional keepers of poultry.

